



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## Large-ring cyclodextrins

Larsen, Kim Lambertsen

*Publication date:*  
2009

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Larsen, K. L. (2009). *Large-ring cyclodextrins*. Abstract from 2° Congresso Nazionale Chimica E Tecnologia delle Ciclodestrine, Asti, Italy.

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

## Large-ring cyclodextrins

Kim Lambertsen Larsen

Section of Chemistry, Department of Biotechnology, Chemistry and Environmental Engineering, Aalborg University, Sohngaardsholmsvej 57, 9000 Aalborg, Denmark

Cyclodextrins are normally considered as small cyclic molecules that, due to a relatively hydrophobic inner cavity, are capable of forming complexes with usually small organic molecules. The only commercially available cyclodextrins (and their derivatives) are made from macrocycles consisting of 6, 7 or 8 glucose units, respectively. These are known as the  $\alpha$ -,  $\beta$ -, and  $\gamma$ -cyclodextrin. However, it is possible to produce cyclodextrins with more glucose units pr. macrocycle. Although, these cyclodextrins may have a degree of polymerization ranging from 9 to several hundreds, they are collectively known as the large-ring cyclodextrins. An overview of the production, purification, structure, properties and possible applications of this highly diverse group of molecules will be given.

